



TITLE:

Positive representations, multiplier Hopf algebra, and continuous canonical basis (String theory, integrable systems and representation theory)

AUTHOR(S):

Ip, Ivan

CITATION:

Ip, Ivan. Positive representations, multiplier Hopf algebra, and continuous canonical basis (String theory, integrable systems and representation theory). 数理解析研究所講究録別冊 2017, B62: 71-86

ISSUE DATE:

2017-05

URL:

<http://hdl.handle.net/2433/243640>

RIGHT:

© 2017 by the Research Institute for Mathematical Sciences, Kyoto University. All rights reserved.

Positive representations, multiplier Hopf algebra, and continuous canonical basis

By

Ivan IP*

Abstract

We introduce the language of multiplier Hopf algebra in the context of positive representations of split real quantum groups, and discuss its applications with a continuous version of Lusztig-Kashiwara’s canonical basis, which may provide a key to prove the closure of the positive representations under tensor products, and harmonic analysis of quantized algebra of functions in the sense of locally compact quantum groups.

§ 1. Introduction

The notion of *positive principal series representations*, or *positive representations* for short, was introduced by I. Frenkel and the author [12] as a new research program devoted to the representation theory of split real quantum groups $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$, where $q = e^{\pi i b^2}$ and its dual $\tilde{q} = e^{\pi i b^{-2}}$. It uses the concept of modular double for quantum groups [7, 8], and has been studied extensively for $U_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$ by Teschner *et al.* [2, 31, 32] in the context of Liouville conformal field theory. Explicit construction of the positive representations \mathcal{P}_{λ} of $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ associated to a simple Lie algebra \mathfrak{g} has been obtained for the simply-laced case in [17] and non-simply-laced case in [18], where the generators $\{E_i, F_i, K_i\}$ of the quantum groups are realized by positive, unbounded, essentially self-adjoint operators acting on $L^2(\mathbb{R}^{l(w_0)})$. Here $l(w_0)$ is the length of the longest element w_0 in the Weyl group W . Furthermore, the so-called *transcendental relations* of the (rescaled) generators:

$$(1.1) \quad \widetilde{\mathbf{e}}_i = \mathbf{e}_i^{\frac{1}{b_i^2}}, \quad \widetilde{\mathbf{f}}_i = \mathbf{f}_i^{\frac{1}{b_i^2}}, \quad \widetilde{K}_i = K_i^{\frac{1}{b_i^2}}$$

Received December 13, 2013. Revised May 28, 2014.

2010 Mathematics Subject Classification(s): Primary 17B37, 20G42, Secondary 47B25

*Department of Mathematics, Faculty of Science, Kyoto University, Kyoto 606-8502, Japan.

give the self-duality between different parts of the modular double, while in the non-simply-laced case, new explicit analytic relations between the quantum group $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ and its Langlands dual $\mathcal{U}_{\tilde{q}}({}^L\mathfrak{g}_{\mathbb{R}})$ have been observed [18].

It becomes clear in the work [15, 19] that the positivity of the generators is immediately connected with the notion of multiplier Hopf algebra introduced by van Daele [35]. Namely, by taking complex powers of the generators as unbounded operators, we can define a C^* -algebraic version of the Drinfeld-Jimbo quantum groups $\mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{g}_{\mathbb{R}})$ that is in some sense dual to certain locally compact quantum groups [24, 25], based on earlier work by Baaĵ and Skandalis [1], Woronowicz [36] and others. This allows us to define the modular double as a quasi-triangular multiplier Hopf algebra with the universal R -operator being the canonical element of the corresponding multiplier Drinfeld's double, giving a braiding structure of the positive representations.

More importantly, we believe that this establishes a link between the C^* -algebraic theory of Connes' non-commutative geometry and Woronowicz's matrix quantum groups, and the theory of Drinfeld-Jimbo type quantum groups at the algebraic and combinatoric level, providing both sides with new machineries.

The construction of $\mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{g}_{\mathbb{R}})$ involves a continuous version of the PBW basis, and subsequently the concept of a continuous version of the *canonical basis*, generalizing the finite dimensional concept first discovered by Lusztig [27, 28], and subsequently proved to exist by Kashiwara [21], using a different method under the name of *global crystal basis*. One can think of it as an analogue to the relation between the (discrete) Fourier series in the harmonic analysis over the circle, which is compact, and the (continuous) Fourier transform in the harmonic analysis over the real line, which is non-compact, with the exponential functions in both cases being the corresponding basis. Furthermore, the use of the remarkable quantum dilogarithm function $G_b(x)$ and its variant $g_b(x)$ introduced earlier by Faddeev and Kashaev [9], will play the role of the q -factorial as well as quantum exponential function in the (continuous) split real setting.

We believe that the nice properties of the canonical basis, especially its behavior under the decomposition of tensor products [21, 29], can be generalized to the continuous version in certain settings, which enables us to complete the proof of the tensor category structure of the category $\text{Rep}^+(\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}}))$ of positive representations.

Moreover, the use of the multiplier Hopf algebra also allows us to study the dual of $\mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{g}_{\mathbb{R}})$, which appears to be the right object for the harmonic analysis of quantized algebra of functions, and coincides with the notion of matrix quantum groups in the locally compact quantum groups setting [24, 25]. With the machinery of the multiplicative unitary, the language of multiplier Hopf algebra provides insight in proving a version of Peter-Weyl theorem for the modular double of split real quantum groups.

As a final remark, the modular double $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$ arising from Liouville theory

is known to be closely related to quantum Teichmüller theory through the construction of conformal blocks [33]. On the other hand, it can also be constructed out of the representation theory of the Borel part of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$, which is the quantum plane [13]. With the help of the continuous canonical basis in the higher rank, the restriction of the positive representations of the modular double to the Borel part $\mathbf{Ub} \subset \mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{g}_{\mathbb{R}})$ is recently studied in [20], which allows us to construct a quantized version of the higher Teichmüller theory as introduced by Fock and Goncharov [11], and connect with Toda field theories [10, 37].

§ 2. Multiplier Hopf Algebra

In this section let us recall the basic definitions of multiplier Hopf algebra. For further details please refer to [35]. One of the original motivation of multiplier Hopf algebra is the following. Consider continuous functions $C[G]$ on a compact group G . Then the multiplication on G induces a coproduct $\Delta : C[G] \longrightarrow C[G] \otimes C[G]$ given by

$$(2.1) \quad \Delta f(g_1, g_2) = f(g_1 g_2).$$

When we discuss locally compact group G , due to the Gelfand-Naimark theorem, it is natural to consider the C^* -algebra $C_0(G)$ of functions vanishing at infinity. However, the formula for Δ above may no longer maps to $C_0(G) \otimes C_0(G)$. Nonetheless, Δf is still bounded and multiplication with elements in $C_0(G) \otimes C_0(G)$ stays in the same space. Hence it is natural to extend the definition of the coproduct and introduce the following notion.

Definition 2.1. Let $\mathcal{B}(\mathcal{H})$ be the algebra of bounded linear operators on a Hilbert space \mathcal{H} . Then the multiplier algebra $M(\mathcal{A})$ of a C^* -algebra $\mathcal{A} \subset \mathcal{B}(\mathcal{H})$ is the C^* -algebra of operators

$$(2.2) \quad M(\mathcal{A}) = \{b \in \mathcal{B}(\mathcal{H}) : b\mathcal{A} \subset \mathcal{A}, \mathcal{A}b \subset \mathcal{A}\}.$$

In particular, \mathcal{A} is an ideal of $M(\mathcal{A})$.

Definition 2.2. A multiplier Hopf $*$ -algebra is a C^* -algebra \mathcal{A} together with the antipode S , the counit ϵ , and the coproduct map

$$(2.3) \quad \Delta : \mathcal{A} \longrightarrow M(\mathcal{A} \otimes \mathcal{A}),$$

all of which can be extended to a map from $M(\mathcal{A})$, such that the usual properties of a Hopf algebra holds on the level of $M(\mathcal{A})$.

In particular, the standard example is $\mathcal{A} = C_0(G)$, and $M(\mathcal{A})$ will be the algebra of bounded functions $C_b(G)$ on the locally compact group G .

Another motivation comes from the definition of the universal R -matrix, which is a very important solution to the Yang-Baxter equation in the theory of compact quantum groups giving the category of finite dimensional representation a braiding structure [5]. However, R is defined as an element of certain completion $\mathcal{U}_h(\mathfrak{g}) \widehat{\otimes} \mathcal{U}_h(\mathfrak{g})$ which is not very natural in the split real setting. In particular, for the theory of modular double $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$, the universal R operator is defined by an integral transformation, and therefore the expression using generators of the split real quantum group does not lie in any algebraic completions. The following definition gives a natural interpretation of R as an element of a multiplier Hopf algebra.

Definition 2.3. A quasi-triangular multiplier Hopf algebra is a multiplier Hopf algebra \mathcal{A} together with an invertible element $R \in M(\mathcal{A} \otimes \mathcal{A})$ such that

$$(2.4) \quad (\Delta \otimes id)(R) = R_{13}R_{23} \in M(\mathcal{A} \otimes \mathcal{A} \otimes \mathcal{A}),$$

$$(2.5) \quad (id \otimes \Delta)(R) = R_{13}R_{12} \in M(\mathcal{A} \otimes \mathcal{A} \otimes \mathcal{A}),$$

$$(2.6) \quad \Delta^{op}(a)R = R\Delta(a) \in M(\mathcal{A} \otimes \mathcal{A}), \quad \forall a \in M(\mathcal{A}),$$

$$(2.7) \quad (\epsilon \otimes id)(R) = (id \otimes \epsilon)(R) = 1 \in M(\mathcal{A}).$$

In Section 4, we will discuss the construction of a C^* -algebraic version of the Drinfeld-Jimbo type quantum group $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ using a continuous version of the canonical basis, such that R naturally arises as the canonical element of this multiplier algebra.

Finally in Section 6, we discuss the application of multiplier Hopf algebra in the context of harmonic analysis on the quantized algebra of functions on $G_{q\tilde{q}}^+(\mathbb{R})$, which is certain dual algebra to $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$. Here the power of multiplier Hopf algebra comes with the tool W called the *multiplicative unitary*. More precisely,

Definition 2.4. Given a GNS-representation of the multiplier Hopf algebra \mathcal{A} on the Hilbert space \mathcal{H} , a *multiplicative unitary* is a unitary operator $W \in \mathcal{B}(\mathcal{H} \otimes \mathcal{H})$ satisfying the pentagon equation

$$(2.8) \quad W_{23}W_{12} = W_{12}W_{13}W_{23},$$

where the leg notation is used.

Then the following key properties are needed:

Proposition 2.5. *The multiplicative unitary encodes the information of the co-product, i.e.*

$$(2.9) \quad W^*(1 \otimes x)W = \Delta(x), \quad x \in \mathcal{A}$$

as operators on $\mathcal{H} \otimes \mathcal{H}$.

Proposition 2.6. *The multiplicative unitary encodes the information of the Hopf dual $\widehat{\mathcal{A}}$, i.e.*

$$(2.10) \quad \widehat{\mathcal{A}} := \{(\omega \otimes 1)W : \omega \in \mathcal{B}(\mathcal{H})^*\}^{norm \text{ closure}} \subset \mathcal{B}(\mathcal{H}),$$

and we have $W \in M(\mathcal{A} \otimes \widehat{\mathcal{A}})$.

Proposition 2.7. *The multiplicative unitary gives the (left) regular co-representation of \mathcal{A} by*

$$(2.11) \quad \Pi : \mathcal{H} \longrightarrow \mathcal{H} \otimes M(\mathcal{A})$$

$$(2.12) \quad f \mapsto W'(f \otimes 1),$$

where W' is W_{21} viewed as an element in $\mathcal{B}(\mathcal{H}) \otimes \mathcal{A}$. By taking the dual we also obtain the (left) regular representation for the dual algebra.

We refer to [15, 34] for more detailed discussions.

§ 3. Positive Representations of $\mathcal{U}_{q\bar{q}}(\mathfrak{g}_{\mathbb{R}})$

The central object in the representation theory of split real quantum groups is the notion of positive representations [12]. In this section we will review its definitions and properties. For simplicity of this exposition, let us only consider simple Lie algebra \mathfrak{g} of simply-laced type [17]. Details for the non-simply-laced case can be found in [18].

Let $q = e^{\pi i b^2}$ with $0 < b < 1$, $b^2 \in \mathbb{R} \setminus \mathbb{Q}$, and consider the usual Drinfeld-Jimbo type quantum group $\mathcal{U}_q(\mathfrak{g})$ generated by $\{E_i, F_i, K_i\}$ for $i \in I$ the root indices, satisfying:

$$(3.1) \quad K_i E_j = q^{a_{ij}} E_j K_i,$$

$$(3.2) \quad K_i F_j = q^{-a_{ij}} F_j K_i,$$

$$(3.3) \quad [E_i, F_j] = \delta_{ij} \frac{K_i - K_i^{-1}}{q - q^{-1}},$$

together with the Serre relations for $i \neq j$:

$$(3.4) \quad E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 = 0,$$

$$(3.5) \quad F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 = 0,$$

where $A := (a_{ij})$ is the Cartan matrix.

Theorem 3.1. [12, 17] *Let*

$$(3.6) \quad e_i := 2 \sin(\pi b^2) E_i, \quad f_i := 2 \sin(\pi b^2) F_i.$$

Note that $2 \sin(\pi b^2) = \left(\frac{i}{q-q^{-1}}\right)^{-1} > 0$. Then there exists a representation \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g})$ parametrized by the \mathbb{R}_+ -span of the cone of positive weights $\lambda \in P_{\mathbb{R}}^+$, or equivalently by $\lambda \in \mathbb{R}_+^n$ where $n = \text{rank}(\mathfrak{g})$, such that

- The generators e_i, f_i, K_i are represented by positive essentially self-adjoint operators acting on $L^2(\mathbb{R}^{l(w_0)})$, where $l(w_0)$ is the length of the longest element $w_0 \in W$ of the Weyl group.
- Define the transcendental generators:

$$(3.7) \quad \tilde{e}_i := e_i^{\frac{1}{b^2}}, \quad \tilde{f}_i := f_i^{\frac{1}{b^2}}, \quad \widetilde{K}_i := K_i^{\frac{1}{b^2}}.$$

Then the representations of the generators $\tilde{e}_i, \tilde{f}_i, \widetilde{K}_i$ are obtained by replacing b with b^{-1} in the representations of the generators e_i, f_i, K_i , and they generate the other part of the modular double $\mathcal{U}_{\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ with $\tilde{q} = e^{\pi i b^{-2}}$.

- The generators e_i, f_i, K_i and $\tilde{e}_i, \tilde{f}_i, \widetilde{K}_i$ commute weakly up to a sign.

The Theorem implies that \mathcal{P}_λ is simultaneously a representation for the split real form $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ and $\mathcal{U}_{\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ by positive operators. Hence we will call \mathcal{P}_λ the positive representation of the modular double

$$\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}}) := \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}}) \otimes \mathcal{U}_{\tilde{q}}(\mathfrak{g}_{\mathbb{R}}).$$

For the non-simply-laced case, the transcendental relations give rise to a direct analytic relation between the quantum group and its Langlands dual [18].

Let us present the expressions of \mathcal{P}_λ in the case of type A_1 and A_2 . By abuse of notation, let us denote by

$$(3.8) \quad [u]e(p) := e^{\pi b(-u+2p)} + e^{\pi b(u+2p)},$$

which is an unbounded positive operator. Here $p := \frac{1}{2\pi i} \frac{\partial}{\partial u}$ is the standard momentum operator.

Proposition 3.2. [2, 32] *The positive representation \mathcal{P}_λ of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$ with $\lambda \in \mathbb{R}_+$ acting on $f(u) \in L^2(\mathbb{R})$ is given by*

$$\begin{aligned} e &= [u - \lambda]e(-p) = e^{\pi b(-u+\lambda-2p)} + e^{\pi b(u-\lambda-2p)}, \\ f &= [-u - \lambda]e(p) = e^{\pi b(u+\lambda+2p)} + e^{\pi b(-u-\lambda+2p)}, \\ K &= e^{-2\pi b u}. \end{aligned}$$

Proposition 3.3. [17] *The positive representation \mathcal{P}_λ of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(3, \mathbb{R}))$ with parameters $\lambda := (\lambda_1, \lambda_2) \in \mathbb{R}_+^2$, corresponding to the reduced expression $w_0 = s_2 s_1 s_2$, acting on $f(u, v, w) \in L^2(\mathbb{R}^3)$, is given by*

$$\begin{aligned} e_1 &= [v - w]e(-p_v) + [u]e(-p_v + p_w - p_u), \\ e_2 &= [w]e(-p_w), \\ f_1 &= [-v + u - 2\lambda_1]e(p_v), \\ f_2 &= [-2u + v - w - 2\lambda_2]e(p_w) + [-u - 2\lambda_2]e(p_u), \\ K_1 &= e^{-\pi b(-u+2v-w+2\lambda_1)}, \\ K_2 &= e^{-\pi b(2u-v+2w+2\lambda_2)}. \end{aligned}$$

§ 4. Continuous Canonical Basis

Again for simplicity, in the following let us consider only the simply-laced case, further discussions can be found in [19]. Recall that in the theory of canonical basis introduced by Lusztig [27, 28], the necessary ingredients are the rescaled simple root generators

$$(4.1) \quad E_i^{(N)} := \frac{E_i^N}{[N]_q!},$$

where $[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}$ is the quantum number, and the non-simple roots vectors given by Lusztig's isomorphism T_i :

$$(4.2) \quad E_{\alpha_k}^{(c)} := T_{i_1} \dots T_{i_{k-1}}(E_{i_k}^{(c)}),$$

such that the canonical basis is given by certain linear combinations of the PBW-basis of the form

$$(4.3) \quad E_{\mathbf{i}}^{\mathbf{c}} := E_{\alpha_N}^{(c_N)} \dots E_{\alpha_2}^{(c_2)} E_{\alpha_1}^{(c_1)},$$

where $w_0 = s_{i_1} \dots s_{i_N}$ is an expression of the longest element of the Weyl group W .

In the context of positive representations, the generators $\{E_i\}_{i \in I}$ of $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ are represented on \mathcal{P}_λ by positive self-adjoint operators. Although the defining formula for Lusztig's isomorphism:

$$(4.4) \quad T_i(E_i) := -F_i K_i^{-1}, \quad T_i(E_j) := q^{\frac{1}{2}} E_j E_i - q^{-\frac{1}{2}} E_i E_j$$

in the compact case does not respect the positivity of the positive representations, it turns out that we can consider instead our favorite rescaled generators $\{\mathbf{e}_i, \mathbf{f}_i\}$ from (3.6), and modify the scaling to obtain

$$(4.5) \quad T_i(\mathbf{e}_i) := q \mathbf{f}_i K_i^{-1}, \quad T_j(\mathbf{e}_i) := \mathbf{e}_{ij} := \frac{q^{\frac{1}{2}} \mathbf{e}_j \mathbf{e}_i - q^{-\frac{1}{2}} \mathbf{e}_i \mathbf{e}_j}{q - q^{-1}},$$

which are again positive self-adjoint, and obey the transcendental relations

$$(4.6) \quad \mathbf{e}_{ij}^{\frac{1}{b^2}} = \widetilde{\mathbf{e}}_{ij} := \frac{\widetilde{q}^{\frac{1}{2}} \widetilde{\mathbf{e}}_j \widetilde{\mathbf{e}}_i - \widetilde{q}^{-\frac{1}{2}} \widetilde{\mathbf{e}}_i \widetilde{\mathbf{e}}_j}{\widetilde{q} - \widetilde{q}^{-1}}.$$

Furthermore, T_i is shown to be represented by conjugation by a unitary element w_i , called the quantum Weyl element. Hence all generators of the form (cf. (4.2))

$$(4.7) \quad \mathbf{e}_{\alpha_k} := T_{i_1} \dots T_{i_{k-1}}(\mathbf{e}_{i_k})$$

are positive self-adjoint and satisfy the transcendental relations. In particular, the expression $\mathbf{e}_{\alpha_k}^{ib^{-1}t}$ is a well-defined bounded unitary operator acting on \mathcal{P}_λ , and invariant under the $b \longleftrightarrow b^{-1}$ duality.

Therefore together with the notation $K_i = q^{H_i}$, we generalize the PBW-basis (4.3) and the following definition is well-defined [19]:

Definition 4.1. Let $n = \text{rank}(\mathfrak{g})$ and $N = l(w_0)$. We define the C^* -algebraic version of the Borel subalgebra

$$\mathbf{Ub} := \mathcal{U}_{q\widetilde{q}}^{C^*}(\mathfrak{b}_{\mathbb{R}}^+)$$

as the operator norm closure of the linear span of all bounded operators on $\mathcal{P}_\lambda = L^2(\mathbb{R}^N)$ of the form

$$(4.8) \quad \vec{F} := F_0(\mathbf{H}) \prod_{k=1}^N \int_C F_k(t_k) \frac{\mathbf{e}_{\alpha_k}^{ib^{-1}t_k}}{G_b(Q + it_k)} dt_k,$$

where $Q = b + b^{-1}$, \mathbf{e}_{α_k} is given by (4.7) and

$$(4.9) \quad F_0(\mathbf{H}) := F_0(ibH_1, \dots, ibH_n)$$

is a smooth compactly supported functions on the positive operators ibH_k , and $F_k(t_k)$ are entire analytic functions that have rapid decay along the real direction. Finally the contour C is along the real axis which goes above the pole of G_b at $t_k = 0$.

Theorem 4.2.

- (1) \mathbf{Ub} is a multiplier Hopf algebra;
- (2) it is invariant under the modular duality $b \longleftrightarrow b^{-1}$;
- (3) it is independent of the choice of the parameter λ ;
- (4) it is independent of the choice of reduced expression of the longest element $w_0 \in W$.

Proof. (1) The quantum dilogarithm function G_b appearing in the definition ensures that \mathbf{Ub} is indeed a multiplier Hopf algebra, where the coproduct of \mathbf{e}_{α_k} will pick up factors of $G_b(Q + it_k)$ due to the q -binomial formula (Lemma A.3) and mutually cancel, giving the correct analytic properties of the bounded operators. We see immediately that $G_b(Q + it)$ plays the role of the q -factorial $[n]_q!$ factors in the compact canonical basis.

(2) The transcendental relations (1.1) and the duality properties (A.3) of G_b ensures that the expression is invariant under $b \longleftrightarrow b^{-1}$, hence it indeed incorporates the properties of the modular double.

(3) From the explicit expression of \mathcal{P}_λ , one notes that the dependence of λ only appears as linear terms in H_i , and the positive representation is injective (see [17]). Hence the algebra does not depend on λ . In fact, there exists a unitary equivalence between $\mathcal{P}_\lambda \simeq \mathcal{P}_{\lambda'}$ for any λ, λ' when restricted to the Borel part, considered in [20].

(4) Finally, we prove in [19] that the continuous span by the basis $\prod_{k=1}^N \frac{\mathbf{e}_{\alpha_k}^{ib^{-1}t_k}}{G_b(Q+it_k)}$ does not depend on the choice of the expression of the longest element w_0 . More precisely, the change of words of w_0 induces an integral transformation on the coefficients $F_k(t_k)$ that preserves their analytic properties. (In the non-simply-laced case however, the explicit formula is not known for type G_2 .) It also shows the existence of the antipode S on the multiplier Hopf algebraic level as an unbounded operator. \square

Now we can apply the construction of Drinfeld's double in the setting of multiplier Hopf algebra [3] to obtain $\mathcal{D} := \mathcal{D}(\mathbf{Ub})$, which is known [4] to be a quasi-triangular multiplier Hopf algebra, where R is given by the canonical element, which is the unique element in $M(\mathcal{D} \otimes \mathcal{D})$ such that

$$(4.10) \quad \langle R, b \otimes a \rangle = \langle a, b \rangle, \quad a \in \mathcal{A}, b \in \mathcal{A}'.$$

Definition 4.3. We define

$$(4.11) \quad \mathbf{U} := \mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{g}_{\mathbb{R}}) := \mathcal{D}(\mathbf{Ub}) / (H'_i = (A^{-1})_{ij}H_j)$$

to be the Drinfeld's double of the Borel subalgebra \mathbf{Ub} modulo the Cartan subalgebra $\mathfrak{h} \subset \mathbf{Ub}^-$.

It follows naturally from the explicit expression of the universal R -operator constructed in [19] which generalizes [22, 26]:

$$(4.12) \quad R = \prod_{ij} q^{\frac{1}{2}(A^{-1})_{ij}H_i \otimes H_j} \prod_{k=1}^N g_b(\mathbf{e}_{\alpha_k} \otimes \mathbf{f}_{\alpha_k}) \prod_{ij} q^{\frac{1}{2}(A^{-1})_{ij}H_i \otimes H_j},$$

together with the integral transformation (A.12) of g_b (replacing the usual q -exponential function), and the Hopf pairing between \mathbf{Ub} and \mathbf{Ub}^- involving $G_b(x)$ (replacing the

usual q -factorial), that the canonical element is given precisely by R . Therefore \mathbf{U} is indeed a quasi-triangular multiplier Hopf algebra. Furthermore, due to Lemma A.6, R induces a unitary transformation between $\mathcal{P}_{\lambda_1} \otimes \mathcal{P}_{\lambda_2} \simeq \mathcal{P}_{\lambda_2} \otimes \mathcal{P}_{\lambda_1}$, hence giving the braiding structure of the positive representations.

Remark 4.4. In the compact case, Lusztig's canonical basis is certain linear combinations of the PBW-basis that is invariant under the bar involution $q \longleftrightarrow q^{-1}$. In the non-compact case, it is expected that the true notion of a continuous canonical basis will be certain integral transformations of the continuous PBW-basis constructed above, which obeys some form of non-compact bar involutions involving complex conjugations. Here the positivity condition is expected to play an important role in its definition.

§ 5. Tensor Products of \mathcal{P}_λ

The notion of continuous canonical basis may provide the key to study the decomposition of tensor products of positive representations. In the finite dimensional situation generated by a highest weight vector ξ_λ with dominant integral weight λ , the $\mathcal{U}_q(\mathfrak{g})$ module is generated by basis of the form $\{E_i^c \xi_\lambda\}$ where E_i^c is given as in (4.3). The decomposition of tensor product is then, roughly speaking, given by the tensor product of the corresponding basis giving the same weight, modulo certain monomial terms with higher order of q in the coefficients [29]. In the setting of positive representations, there are no highest weight vector anymore, but still one can fix an initial weight vector ξ_λ and define \mathcal{P}_λ to be “generated” by generalized functions such as

$$(5.1) \quad \mathcal{P}_\lambda = \text{“continuous span”} \left\{ \prod_{k=1}^N \frac{\mathbf{e}_{\alpha_k}^{ib^{-1}t_k}}{G_b(Q + it_k)} \xi_\lambda \right\}.$$

As an example, in the case of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$, one can regard $\mathcal{P}_\lambda = L^2(\mathbb{R})$ as generated by the initial weight vector $\delta(x)$, (where K acts as $e^{-2\pi b\lambda}$), and look at the action of $\mathbf{e}^{ib^{-1}t}$ on $\delta(x)$ [19]:

$$(5.2) \quad \mathbf{e}^{ib^{-1}t} \cdot \delta(x) = e^{\frac{\pi i(t-2\lambda)t}{2}} \frac{G_b(\frac{Q}{2} - i\lambda + it)}{G_b(\frac{Q}{2} - i\lambda)} \delta(x - t).$$

In particular, in the analytic continuation $t \longrightarrow -ib$, we recover the action of \mathbf{e} given in Proposition 3.2.

One can then take these continuous basis and follow the decomposition in the compact case, generalizing the formula for the decomposition $\mathcal{P}_{\lambda_1} \otimes \mathcal{P}_{\lambda_2}$. In principle this will recover the intertwiner:

$$(5.3) \quad \mathcal{P}_{\lambda_1} \otimes \mathcal{P}_{\lambda_2} \simeq \int_{\mathbb{R}_+}^{\oplus} \mathcal{P}_\lambda \sinh(2\pi b\lambda) \sinh(2\pi b^{-1}\lambda) d\lambda$$

given by a series of integral transformations involving $G_b(x)$ proved in [30, 32].

Of course, one has to be careful when dealing with such generalized functions in terms of functional analysis. Nevertheless, investigating this observation, we have the following more precise conjecture for the higher rank case.

Conjecture 5.1. *The positive representations for $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ is closed under taking the tensor product, and it decomposes as*

$$(5.4) \quad \mathcal{P}_{\alpha} \otimes \mathcal{P}_{\beta} \simeq \int_{\mathbb{R}_+^N}^{\oplus} \mathcal{P}_{\vec{\gamma}} d\mu(\vec{\gamma}),$$

where $\vec{\gamma} = \sum_{\alpha \in \Delta_+} \gamma_{\alpha} \omega_{\alpha}$ suming over all the positive roots, where $\gamma_{\alpha} \in \mathbb{R}_+$ and ω_{α} are the fundamental weights, with the abuse of notation $\omega_{\alpha} := \omega_{\alpha_1} + \omega_{\alpha_2}$ if $\alpha := \alpha_1 + \alpha_2$ is not simple. The Plancherel measure is given by

$$d\mu(\vec{\gamma}) = \prod_{\alpha \in \Delta_+} \sinh(2\pi b \gamma_{\alpha}) \sinh(2\pi b^{-1} \gamma_{\alpha}) d\gamma_{\alpha}.$$

Remark 5.2. It was pointed out by Masahito Yamazaki that this Plancherel measure naturally comes from a somewhat indirect connection from supersymmetric gauge theory, which is discussed for example in [6, 14].

In particular, in the simply-laced case, we believe it suffices to prove

Conjecture 5.3. *The tensor product of positive representations for $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(3, \mathbb{R}))$ decomposes as*

$$(5.5) \quad \mathcal{P}_{(\alpha_1, \alpha_2)} \otimes \mathcal{P}_{(\beta_1, \beta_2)} \simeq \int_{\mathbb{R}_+^3}^{\oplus} \mathcal{P}_{(\gamma_1 + \gamma_3, \gamma_2 + \gamma_3)} d\mu(\vec{\gamma}).$$

Finally, let us remark that the situation is simpler if we restrict the positive representations \mathcal{P}_{λ} to the Borel part $\mathfrak{b}_{\mathbb{R}} \subset \mathfrak{g}_{\mathbb{R}}$. Using the notion of $\mathcal{U}_{q\tilde{q}}^{C^*}(\mathfrak{b}_{\mathbb{R}})$ and the theory of multiplicative unitary associated to its GNS-representations, it is shown in a recent study [20] that the positive representations \mathcal{P}_{λ} restricted to $\mathfrak{b}_{\mathbb{R}}$ does not depend on λ , and it is closed under taking tensor products. This construction induces the quantum mutation operator \mathbf{T} which will be a key ingredient to obtain a candidate for the quantum higher Teichmüller theory, generalizing earlier work by Frenkel-Kim [13].

§ 6. Harmonic Analysis of $L^2(G_{q\tilde{q}}^+(\mathbb{R}))$

In the last section, we would like to discuss the application of multiplier Hopf algebra in the context of harmonic analysis on the quantized algebra of functions on

$G_{q\tilde{q}}^+(\mathbb{R})$, which is certain dual algebra to $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$. This is motivated from the following form of Peter-Weyl theorem [15, 31]

$$(6.1) \quad L^2(SL_{q\tilde{q}}^+(2, \mathbb{R})) \simeq \int_{\mathbb{R}_+}^{\oplus} \mathcal{P}_{\lambda} \otimes \mathcal{P}_{\lambda}^* d\mu(\lambda)$$

as left and right regular representations of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$, where the Plancherel measure $d\mu(\lambda)$ is the same as before.

Here, the Hilbert space $L^2(SL_{q\tilde{q}}^+(2, \mathbb{R}))$ is constructed as follows. Starting with the *quantum plane* generated by positive self-adjoint elements $\{A, B\}$ with $AB = q^2 BA$, one define the C^* -algebraic version $\mathcal{A}_{q\tilde{q}}$ similar to the construction of \mathbf{Ub} , giving it a multiplier Hopf algebra structure, as well as imposing the modular duality $b \longleftrightarrow b^{-1}$. With the existence of a Haar functional and certain density conditions, $\mathcal{A}_{q\tilde{q}}$ can be shown to be a locally compact quantum group in the sense of [24, 25]. Finally, by applying Woronowicz's *quantum double construction*, which is the dual of Drinfeld's double construction, we obtain a multiplier Hopf algebra $\mathcal{A}(SL_{q\tilde{q}}^+(2, \mathbb{R}))$, and the Hilbert space $L^2(SL_{q\tilde{q}}^+(2, \mathbb{R}))$ is defined by its von Neumann completion using the GNS representation. The analysis utilizes the power of the multiplicative unitary W associated to the GNS representation of the quantum plane, and the corresponding quantum double counterparts, which is the key ingredient to construct the regular representations of $\mathcal{U}_{q\tilde{q}}(\mathfrak{sl}(2, \mathbb{R}))$.

One natural question is the extension of this result to the higher rank. In [16], we define the *Gauss-Lusztig's decomposition* (for type A_n), which plays the role of Woronowicz's quantum double construction. Here using Lusztig's data for the totally positive matrices, the space $G_{q\tilde{q}}^+(\mathbb{R})$ is given by an algebra generated by elements that form interconnecting q -tori. Different parametrization of the Lusztig's data gives intertwiners that is also closely related to the positive representations on a combinatorial level [17, 23]. Imposing positivity, we can mimic the construction in the case of the quantum plane and give this algebra a multiplier Hopf algebra structure, as well as its von Neumann completion, hence the Hilbert space $L^2(G_{q\tilde{q}}^+(\mathbb{R}))$. One can then discuss the regular representations of $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$ acting on this space, and give the following conjecture.

Conjecture 6.1. *We have the decomposition*

$$(6.2) \quad L^2(G_{q\tilde{q}}^+(\mathbb{R})) \simeq \int_{\mathbb{R}_+^n}^{\oplus} \mathcal{P}_{\lambda} \otimes \mathcal{P}_{\lambda}^* d\mu(\lambda)$$

as left and right regular representations of $\mathcal{U}_{q\tilde{q}}(\mathfrak{g}_{\mathbb{R}})$. This time, however, λ runs through the fundamental weights ω_i corresponding to simple roots only.

§ A. Quantum Dilogarithms

First introduced by Faddeev and Kashaev [9], the quantum dilogarithm $G_b(x)$ and its variant $g_b(x)$ play a crucial role in the study of positive representations of split real quantum groups, and also appear in many other areas of mathematics and physics. In this appendix, let us recall the definition and some properties of the quantum dilogarithm functions [2, 15, 32] that is needed in this paper.

Definition A.1. The quantum dilogarithm function $G_b(x)$ is defined on $0 \leq \operatorname{Re}(z) \leq Q$ by

$$(A.1) \quad G_b(x) = \bar{\zeta}_b \exp \left(- \int_{\Omega} \frac{e^{\pi t z}}{(e^{\pi b t} - 1)(e^{\pi b^{-1} t} - 1)} \frac{dt}{t} \right),$$

where

$$(A.2) \quad \zeta_b = e^{\frac{\pi i}{2}(\frac{b^2+b^{-2}}{6} + \frac{1}{2})},$$

and the contour goes along \mathbb{R} with a small semicircle going above the pole at $t = 0$. This can be extended meromorphically to the whole complex plane with poles at $x = -nb - mb^{-1}$ and zeros at $x = Q + nb + mb^{-1}$, for $n, m \in \mathbb{Z}_{\geq 0}$;

The quantum dilogarithm $G_b(x)$ satisfies the following properties:

Proposition A.2. *Self-duality:*

$$(A.3) \quad G_b(x) = G_{b^{-1}}(x);$$

Functional equations:

$$(A.4) \quad G_b(x + b^{\pm 1}) = (1 - e^{2\pi i b^{\pm 1} x}) G_b(x);$$

Reflection property:

$$(A.5) \quad G_b(x) G_b(Q - x) = e^{\pi i x(x - Q)};$$

Complex conjugation:

$$(A.6) \quad \overline{G_b(x)} = \frac{1}{G_b(Q - \bar{x})},$$

in particular

$$(A.7) \quad \left| G_b\left(\frac{Q}{2} + ix\right) \right| = 1 \text{ for } x \in \mathbb{R};$$

Asymptotic properties:

$$(A.8) \quad G_b(x) \sim \begin{cases} \bar{\zeta}_b & \operatorname{Im}(x) \rightarrow +\infty \\ \zeta_b e^{\pi i x(x - Q)} & \operatorname{Im}(x) \rightarrow -\infty \end{cases},$$

Lemma A.3 (*q*-binomial theorem). *For positive self-adjoint variables U, V with $UV = q^2VU$, we have:*

$$(A.9) \quad (V + U)^{ib^{-1}t} = \int_C \begin{pmatrix} it \\ i\tau \end{pmatrix}^b V^{ib^{-1}(t-\tau)} U^{ib^{-1}\tau} d\tau,$$

where the *q*-beta function (or *q*-binomial coefficient) is given by

$$(A.10) \quad \begin{pmatrix} t \\ \tau \end{pmatrix}^b = \frac{G_b(Q + t)}{G_b(Q + \tau)G_b(Q + t - \tau)},$$

and C is the contour along \mathbb{R} that goes above the pole at $\tau = 0$ and below the pole at $\tau = t$.

Definition A.4. The function $g_b(x)$ is defined by

$$(A.11) \quad g_b(x) = \frac{\overline{\zeta_b}}{G_b(\frac{Q}{2} + \frac{\log x}{2\pi ib})},$$

where \log takes the principal branch of x .

Lemma A.5. *[2, (3.31), (3.32)] We have the following Fourier transformation formula:*

$$(A.12) \quad \int_{\mathbb{R}+i0} \frac{e^{-\pi it^2}}{G_b(Q + it)} X^{ib^{-1}t} dt = g_b(X),$$

where X is a positive operator and the contour goes above the pole at $t = 0$.

Lemma A.6. *By (A.7), $|g_b(x)| = 1$ when $x \in \mathbb{R}_+$, hence $g_b(X)$ is a unitary operator for any positive operator X . Furthermore we have the self-duality of $g_b(x)$ given by*

$$(A.13) \quad g_b(X) = g_{b^{-1}}(X^{\frac{1}{b^2}}).$$

Acknowledgments

I would like to thank the organizing committee of the RIMS symposium on String Theory, Integrable Systems and Representation Theory for their invitation. I would also like to thank in particular Hiraku Nakajima, Masato Okado, Tomoki Nakanishi and Yoshiyuki Kimura for stimulating interesting discussions and providing useful comments. This work was supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan.

References

- [1] S. Baaĵ, G. Skandalis, *Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres*. In *Annales scientifiques de l'Ecole normale supérieure*, Société mathématique de France **26** (4), 425-488, (1993)
- [2] A.G. Bytsko, K. Teschner, *R-operator, co-product and Haar-measure for the modular double of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$* , Comm. Math. Phys., **240**, 171-196, (2003)
- [3] L. Delvaux, A. van Daele, *The Drinfel'd double of multiplier Hopf algebras*, Jour. Alg., **272**(1), 273-291, (2004)
- [4] L. Delvaux, A. van Daele, *The Drinfel'd double versus the Heisenberg double for an algebraic quantum group*, Jour. Pure App. Alg., **190**, Issues 1-3, 59-84, (2004)
- [5] V. G. Drinfeld, *Hopf algebras and the quantum Yang-Baxter equation*, Doklady Akademii Nauk SSSR, **283** (5), 1060-1064, (1985)
- [6] N. Drukker, D. Gaiotto, and J. Gomis, *The Virtue of Defects in 4D Gauge Theories and 2D CFTs*, Journal of High Energy Physics, **6**, 1-54, (2011)
- [7] L.D. Faddeev, *Discrete Heisenberg-Weyl group and modular group*, Lett. Math. Phys., **34**, 249-254, (1995)
- [8] L.D. Faddeev, *Modular double of quantum group*, arXiv:math/9912078v1 [math.QA], (1999)
- [9] L.D. Faddeev, R.M. Kashaev, *Quantum dilogarithm*, Modern Phys. Lett. **A9**, 427-434, (1994)
- [10] V. A. Fateev, A. V. Litvinov, *Correlation functions in conformal Toda field theory I*, Journal of High Energy Physics, **11**, 002, (2007)
- [11] V. Fock, A. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publications Mathématiques de l'Institut des Hautes Études Scientifiques, **103** (1), 1-211, (2006)
- [12] I. Frenkel, I. Ip, *Positive representations of split real quantum groups and future perspectives*, Int. Math. Res. Notices, Vol 2014 (8), 2126-2164, (2014)
- [13] I. Frenkel, H. Kim, *Quantum Teichmüller space from quantum plane*, Duke Math. J., **161** (2), 305-366, (2012)
- [14] K. Hosomichi, S. Lee, J. Park, *AGT on the S-duality Wall*, Journal of High Energy Physics, **12**, 1-15, (2010)
- [15] I. Ip, *Representation of the quantum plane, its quantum double and harmonic analysis on $GL_q^+(2, \mathbb{R})$* , Selecta Mathematica, **19** (4), 987-1082, (2013)
- [16] I. Ip, *Gauss-Lusztig decomposition of $GL_q^+(n, \mathbb{R})$ and representations by q -tori*, Journal of Pure and Applied Algebra, Vol 219 (12), 5650-5672, (2015)
- [17] I. Ip, *Positive representations of split real simply-laced quantum groups*, arXiv:1203:2018, (2012)
- [18] I. Ip, *Positive representations of non-simply-laced split real quantum groups*, Journal of Algebra, Vol 425, 245-276, (2015)
- [19] I. Ip, *Positive Representations of Split Real Quantum Groups: The Universal R Operator*, International Mathematics Research Notices, Vol 2015 (1), 240-287, (2015)
- [20] I. Ip, *On tensor products of positive representations of split real quantum Borel subalgebra $\mathcal{U}_{q\bar{q}}(\mathfrak{b}_{\mathbb{R}})$* , Transactions of the American Mathematical Society, doi:10.1090/tran/7110 (to appear) (2016)
- [21] M. Kashiwara, *On Crystal Bases of the Q -Analogue of Universal Enveloping Algebras*, Duke J., Vol 63 (2), (1991), 465-516

- [22] A. Kirillov, N. Reshetikhin, *q-Weyl group and a multiplicative formula for universal R matrices*, Commun. Math. Phys., **134**, 421-431, (1990)
- [23] A. Kuniba, M. Okado, Y. Yamada, *A Common Structure in PBW Bases of the Nilpotent Subalgebra of $U_q(\mathfrak{g})$ and Quantized Algebra of Functions*, SIGMA **9**, 049 (2013)
- [24] J. Kustermans and S. Vaes, *Locally compact quantum groups*, Ann. Sci. Ecole Norm. Sup. (4) **33**, 837-934, (2000)
- [25] J. Kustermans and S. Vaes, *Locally compact quantum groups in the von Neumann algebraic setting*, Math. Scand. **92**, 68-92, (2003)
- [26] S. Levendorskiĭ, Ya. Soibelman, *Some applications of the quantum Weyl groups*, Jour. Geom. Phys., **7** No. 2, 241-254, (1990)
- [27] G. Lusztig, *Canonical Bases Arising from Quantized Enveloping Algebras*, Jour. AMS, Vol 3., No. 2 (1990), 447-498
- [28] G. Lusztig, *Canonical Bases Arising from Quantized Enveloping Algebras. II*, Prog. Theo. Phys. Supp. No.102, (1990), 175-201
- [29] G. Lusztig, *Canonical bases in tensor products*, Proc. Natl. Acad. Sci. USA, Vol 89, (1992), 8177-8179
- [30] I. Nidaiev, J. Teschner *On the relation between the modular double of $U_q(\mathfrak{sl}(2, \mathbb{R}))$ and the quantum Teichmüller theory*, arXiv:1302.3454 (2013)
- [31] B. Ponsot, J. Teschner, *Liouville bootstrap via harmonic analysis on a noncompact quantum group*, arXiv: hep-th/9911110, (1999)
- [32] B. Ponsot, J. Teschner, *Clebsch-Gordan and Racah-Wigner coefficients for a continuous series of representations of $U_q(\mathfrak{sl}(2, \mathbb{R}))$* , Comm. Math. Phys., **224**, 613-655, (2001)
- [33] J. Teschner, *On the relation between quantum Liouville theory and the quantized Teichmüller spaces*, Int. J. Modern Phys A **19** (supp02), 459-477, (2004)
- [34] T. Timmermann, *An invitation to quantum groups and duality*, EMS Text-books in Mathematics, (2008)
- [35] A. van Daele, *Multiplier Hopf algebras*, Trans. Amer. Math. Soc. **342** 917-932, (1994)
- [36] S.L. Woronowicz, *From multiplicative unitaries to quantum groups*, Int. J. Math. **7**(1), 127-149, (1996)
- [37] N. Wyllard, *A_{N-1} conformal Toda field theory correlation functions from conformal $\mathcal{N} = 2$ $SU(N)$ quiver gauge theories*, Journal of High Energy Physics, **11** 002, (2009)